

Iron Standard, Atomic Absorption

48937-160	100 mL	18.12
48937-320	500 mL	46.75

Actual Assay on the label1000 µg/mL Fe (17.91 mmol. l⁻¹)In dilute HNO₃

Traceable to NIST standards

Isonicotinic Acid 99%

C ₆ H ₅ NO ₂	FW 123.11
CAS 55-22-1	

49266-140	100 g	79.90
-----------	-------	-------

Melting point310 - 315° C (subl.)

Isopropyl Ether, Reagent, A.C.S.

(C ₃ H ₇) ₂ O	FW 102.18
CAS 108-20-3	

49375-360	1L	80.30
49375-540	4L	180.41

A.C.S. Specifications

Assay [(C ₃ H ₇) ₂ O]99.0% min.
Color (APHA)25 max.
Peroxide (as/en C ₆ H ₁₄ O ₂)05% max.
Residue after evaporation01% max.
Titration acid00007 meq/g max.

Isoquinoline, 97%

C ₉ H ₇ N	FW 129.15
CAS 119-65-3	

49404-140	100 g	150.72
-----------	-------	--------

Melting point26 - 28° C

Karl Fischer Reagent, Stabilized**Single solution**

49634-320	500 mL	76.32
49634-360	1 L	125.08
49634-540	4 L	309.52

Karl Fischer - Solvent Mixture Diluent

(2-Methoxyethanol and Pyridine

For adjusting Karl Fischer Reagent

49646-360	1 L	129.32
-----------	-----	--------

Lactic Acid, 85%, Reagent, A.C.S.

CH ₃ CHOHCOOH	FW 90.08
CAS 50-21-5	

49864-320	500 mL	65.34	6X500 mL	313.48
49864-400	2 L	108.58	6X2 L	521.01
49864-540	4 L	184.23	4X4 L	626.38

A.C.S. Specifications

Assay (C ₃ H ₆ O ₃)85.0 - 90.0%
Substances darkened by Sulfuric AcidPT.
Residue after ignition02% max.
Chloride(Cl)001% max.
Sulfate (SO ₄)002% max.
Heavy metals (as Pb)5 ppm max.
Iron (Fe)5 ppm max.

Lactic Acid, Volumetric Solution, 1 N

49883-360	1 L	49.09
-----------	-----	-------

1mL = 90.08mg C₃H₆O₃

Normality0.995 - 1.005

Traceable to NIST standards

Lactose, Powder, Lab-Grade

C ₁₂ H ₂₂ O ₁₁ • H ₂ O	FW 360.32
CAS 64044-51-5	

49968-300	500 g	36.65	6X500 g	197.96
-----------	-------	-------	---------	--------

Lanolin, Anhydrous, Lab-Grade

CAS 8006-54-0

50048-580	10 kg	223.06
-----------	-------	--------