

Lanthanum Chloride, Hydrated, Reagent, A.C.S.

LaCl₃ • (6-7)H₂O FW 245.27 + xH₂O
 CAS 10099-58-8

50082-080	25 g	27.35
50082-140	100 g	73.74
50082-300	500 g	472.02

A.C.S. Specifications

Assay (LaCl ₃)	.64.5 - 70.0%
Insoluble matter	.01% max.
Calcium (Ca)	.10 ppm max.
Magnesium (Mg)	.10 ppm max.

Lanthanum Oxide, 99.99%

La₂O₃ FW 325.81
 CAS 1312-81-8

50186-140	100 g	115.43
50186-300	500 g	233.07

Lead, Granular, Reagent

Pb AW 207.20
 CAS 7439-92-1

50370-300	500 g	64.98	6X500 g	312.09
-----------	-------	-------	---------	--------

Specifications

Assay (Pb)	.99.5% min.
Antimony and Tin (as Sn)	.005% max.
Arsenic (As)	.1 ppm max.
Bismuth (Bi)	.5 ppm max.
Copper (Cu)	.3 ppm max.
Iron (Fe)	.001% max.
Nickel (Ni)	.001% max.
Silver (Ag)	.2 ppm max.

Lead Standard, Atomic Absorption

50358-160	100 mL	18.12
50358-320	500 mL	46.75

1000 µg/mL Pb (4.83 mmol.l⁻¹)
 In dilute HNO₃

Traceable to NIST standards
 Actual Assay on the label

Lead, Test, Granular

Suitable for use in fire assay.

Pb FW 207.20
 CAS 7439-92-1

029-761-01	50 lb	244.65
------------	-------	--------

Lead Acetate, Crystals, Reagent, A.C.S.

(CH₃COO)₂Pb • 3H₂O FW 379.30
 CAS 6080-56-4

50600-300	500 g	79.93	6X500 g	384.27
50600-380	2 kg	196.69	6X2 kg	943.82
50600-580	10 kg	570.53		
50600-820	50 kg	2564.90		

A.C.S. Specifications

Assay [(CH ₃ COO) ₂ Pb • 3H ₂ O]	.99.0 - 103.0%
Insoluble matter	.01% max.
Chloride (Cl)	.5 ppm max.
Nitrate & Nitrite (as NO ₃)	.005% max.
Calcium (Ca)	.005% max.
Copper (Cu)	.002% max.
Iron (Fe)	.001% max.
Potassium (K)	.005% max.
Sodium (Na)	.01% max.

Lead Carbonate, Basic, Powder, Reagent, A.C.S.

(PbCO₃)₂ • Pb(OH)₂ FW 775.67
 CAS 1319-46-6

50692-140	100 g	56.79
-----------	-------	-------

A.C.S. Specifications

Insoluble in dilute acetic acid	.02% max.
Chloride (Cl)	.002% max.
Nitrate & Nitrite (as NO ₃)	.P.T.
Cadmium (Cd)	.002% max.
Calcium (Ca)	.01% max.
Iron (Fe)	.005% max.
Potassium (K)	.02% max.
Sodium (Na)	.05% max.
Zinc (Zn)	.003% max.