

2-Mercaptoethanol, 98%

HSCH₂CH₂OH FW 78.14
CAS 60-24-2

M370-1-100g 100 g 61.73
M370-1-500g 500 g 132.00

Boiling point157° C

Mercuric Acetate, Powder, Reagent, A.C.S.

(CH₃COO)₂Hg FW 318.68
CAS 1600-27-7

56074-140 100 g 108.81

A.C.S. Specifications

Assay [(CH₃COO)₂Hg]98.0% min.
Insoluble matter0.01% max.
Residue after reduction0.02% max.
Chloride (Cl)0.005% max.
Nitrate (NO₃)0.005% max.
Sulfate (SO₄)0.005% max.
Other heavy metals (as Pb)0.002% max.
Iron/Fer (Fe)0.001% max.
Mercurous mercury (as Hg)0.4% max.

Mercuric Chloride, Crystals, Reagent, A.C.S.

HgCl₂ FW 271.50
CAS 7487-94-7

56166-140 100 g 65.72
56166-300 500 g 252.28 6X500 g 1210.94
56166-380 2 kg 685.29 6X2 kg 3289.18

A.C.S. Specifications

Assay (HgCl₂)99.5% min.
Solution in etherP.T.
Residue after reduction0.02% max.
Iron(Fe)0.002% max.

Mercuric Chloride 0.1 M Solution

C5062 1L 49.79
C5062 4L 119.44
C5062 20L 298.71

Mercuric Iodide, Red, Powder, Reagent, A.C.S.

HgI₂ FW 454.40
CAS 7774-29-0

56258-140 100 g 75.85

A.C.S. Specifications

Assay (dried basis) (HgI₂)99.0% min.
Solubility in KI solutionP.T.
Mercurous mercury (as Hg)0.1% max.
Soluble mercury salts (as Hg)0.05% max.

Mercuric Nitrate, Crystals, Reagent, A.C.S.

Hg(NO₃)₂ • H₂O FW 342.62
CAS 7783-34-8

53338-080 25 g 40.44
53338-140 100 g 84.61

A.C.S. Specifications

Assay [Hg(NO₃)₂ • H₂O]98.0% min.
Residue after reduction0.01% max.
Chloride (Cl)0.002% max.
Sulfate (SO₄)0.002% max.
Iron (Fe)0.001% max.

Mercuric Nitrate, 0.141 N, Acculute

Concentrate to prepare 1 L
56350-000 6X amp 154.29

Mercuric Nitrate, 0.0141 N, Acculute

Concentrate to prepare 1 L
56357-000 6X amp 89.80

Mercuric Nitrate, 0.01 N, Acculute

Concentrate to prepare 1 L
56354-000 6X amp 134.58