

Molecular Sieves, 4A, Mesh 4 - 8

60674-300 500 g 47.38

Molybdate Reagent

60778-540 4 L 104.44

Molybdenum Standard, Atomic Absorption

60805-160 100 mL 18.12

60805-320 500 mL 46.75

Actual Assay on the label

1000 µg/mL Mo (10.42 mmol. l⁻¹)

In dilute NH₄OH

Traceable to NIST standards

Morpholine, Reagent, A.C.S.

C₄H₉NO FW 87.12
CAS 110-91-8

61076-320 500 mL 47.09

61076-360 1 L 88.51

A.C.S. Specifications

Assay (C₄H₉NO)99.0% min.

Color (APHA)15 max.

Boiling range126.00-130.0°C

Murexide, Anhydrous, Reagent, A.C.S.

C₈H₈N₆O₆ FW 284.20
CAS 3051-09-0

61180-040 5 g 31.61

61180-300 500 g 130.38

A.C.S. Specifications

Suitability as a complexometric indicatorP.T.

Naphthalene, Recrystallized

C₁₀H₈ FW 128.17
CAS 91-20-3

61318-300 500 g 43.25

Melting range79 - 81° C

α-Naphtholphthalein, Indicator

C₂₈H₁₈O₄ FW 418.45
CAS 596-01-0

61794-060 10 g 431.88

pH range:7.0-9.0

N-(1-Naphthyl)-ethylenediamine, Dihydrochloride, A.C.S.

C₁₂H₁₄N₂ • 2HCl FW 259.18
CAS 1465-25-4

61870-060 10 g 35.18

61870-080 25 g 69.77

A.C.S. Specifications

Sensitivity to sulfanilamideP.T.

SolubilityP.T.

Water(H₂O)5% max.

Nessler Reagent, Solution

61930-320 500 mL 57.24

61930-360 1 L 83.74

Nickel Standard, Atomic Absorption

61957-160 100 mL 18.12

61957-320 500 mL 46.75

Actual Assay on the label

1000 µg/mL Ni (17.04 mmol. l⁻¹)

In dilute HNO₃

Traceable to NIST standards